MULTIMEDIA UNIVERSITY

STUDENT ID NO

MULTIMEDIA UNIVERSITY

FINAL EXAMINATION

TRIMESTER 2, 2019/2020

BMS1824 – MANAGERIAL STATISTICS

(All sections / Groups)

06 MARCH 2020
9.00 a.m. - 11.00 a.m.
(2 Hours)

INSTRUCTIONS TO STUDENTS

1. This question paper consists of **TWELVE (12)** printed pages with:
   Section A: Ten (10) multiple choice questions (20%)
   Section B: Three (3) structured questions (80%)

2. Answer **ALL** questions.

3. Answer **Section A** in the multiple-choice answer sheet provided and **Section B** in the answer booklet provided.

4. Students are allowed to use non-programmable scientific calculators with no restrictions.

SECTION A: MULTIPLE CHOICE QUESTIONS (20 MARKS)

There are TEN (10) questions in this section. Answer ALL questions on the multiple choice answer sheet.

1. Questionnaire survey method is used to collect _____.

   A. Secondary data
   B. Qualitative variable
   C. Primary data
   D. Discrete data

2. Which of these represent qualitative data?

   A. Height of a student.
   B. Response on liking or disliking of a product.
   C. The income of a government servant in a city.
   D. The number of students in a class.

3. The middle value of an ordered array of numbers is the _____.

   A. Mode
   B. Mean
   C. Median
   D. Standard deviation

4. Which branch of statistics deals with the techniques that are used to organize, summarize and present the data?

   A. Advanced statistics
   B. Probability statistics
   C. Inferential statistics
   D. Descriptive statistics

5. The coefficient of skewness is always zero for _____.

   A. Symmetrical dataset.
   B. Positively skewed
   C. Negatively skewed
   D. None of these

Continued...

6. Determine which statement is TRUE?

A. A single, extremely large value can affect the median more than the mean.
B. In a data set, the mode will always be unique.
C. The height of basketball players is considered as a continuous variable.
D. The positive square root of the variance is called as the coefficient of variation.

7. If the standard deviation of a population is 9, the population variance is:

A. 9.
B. 3.
C. 21.
D. 81.

8. Consider the following data:

14 23 9 12 21 18 8

What is the range?

A. 15
B. 9 to 21
C. 14
D. 8

9. Find the median of the following data:

180 280 320 160 200 300 400

A. 140
B. 300
C. 180
D. 280

10. The mean of a distribution is 23, the median is 24 and the mode is 25.5. It is most likely that this distribution is _____.

A. Left skewed
B. Symmetrical
C. Asymptotic
D. Right skewed

Continued...

SECTION B: STRUCTURED QUESTIONS (80 Marks)

There are THREE questions in this section. Candidates MUST answer ALL THREE questions.

Question 1 (25 Marks)

a) A restaurant chain routinely surveys its customers whether he or she would return to the restaurant and also to rate on the quality of food. From 300 customers, it recorded the following:

Rating on Food Quality	Will Return	Will not return
Poor	6	30
Good	129	69
Excellent	60	6

i. What is the probability that the customers rate the food as good? [3 marks]

ii. What is the probability that the customer rate the food as excellent and will return to the restaurant? [3 marks]

iii. Given that the customer rate the food as good, what is the probability that he or she will return to the restaurant? [4 marks]

b) A concerned parents group determined the number of commercials shown in each of five children's programs within one hour of the TV programs as follows:

Number of Commercials, X	5	6	7	8	9
Probability, P(X)	0.2	0.25	0.38	0.10	0.07

i. What would be the average number of commercials over one hour of the children's programs? [4 marks]

ii. Compute the standard deviation. [5 marks]

c) A survey from Teenage Research Group in Illinois found that 30 percent of teenage consumers receive their spending money from part-time jobs. If 5 teenagers are selected at random, find the probability that at least 3 of them will have part-time jobs. [6 marks]

Continued...

Question 2 (25 Marks)

a) A survey of 15 large cities in US finds that the average of one-way commute time is 25.4 minutes or more. A chamber of commerce executive feels that the commute time is less than that and wants to publicize this. He randomly selects 25 commuters and found that the average commute time is 22.1 minutes with a standard deviation of 5.3 minutes.
At α=0.10, is there enough evidence to support his claim? [13 marks]

b) A researcher wishes to estimate the number of days it takes by an automobile dealer to sell a Honda Jazz. A sample of 50 cars had a mean time on the dealer's job log of 54 days. Assume the population standard deviation to be 6.0 days.

i. Construct a 95% confidence interval for the average number of days taken by the dealer to sell a Honda Jazz. [6 marks]

ii. The researcher wishes to conduct another survey on the above. How large a sample is needed to estimate the mean number of days to within 2 days, with 90 percent confidence? [6 marks]

Question 3 (30 Marks)

a) A finance manager wants to determine the relationship between the interest rates for car loans and the average number of customers who apply for the loans in a month from a finance company. The data for the sample of 12 months are regressed and the summary output of the relationship between the two variables are shown below:



Continued....

SUMMARY OUTPUT

Regression Statistics	
Multiple R	- 0.985931
R Square	0.972061
Adjusted R Square	0.969267
Standard Error	2.538413
Observations	12

ANOVA

	df	*SS*	*MS*	*F*	*Significance F*
Regression	1	2241.815	2241.815	347.9167	0.0000049
Residual	10	64.43539	6.443539		
Total	11	2306.25			

	Coefficients	*Standard Error*	*t Stat*	*P-value*
Intercept	180.4905	6.407955	28.16662	0.0000000
Interest Rate (%)	-16.1369	0.865131	-18.6525	0.0007

i. Based on the scatter plot, describe the relationship between interest rate and number of car loan applicants. [3 marks]

ii. State the estimated linear regression equation for the above model. [3 marks]

iii. What is the correlation coefficient for the regression model? Interpret the value. [3 marks]

iv. What is the coefficient of determination for the regression model? Interpret the value. [4 marks]

v. At 5% significance level, is there any significant relationship between interest rate and number of car loan applicants? Use p-value approach. [4 marks]

vi. Predict the number of applicant if the interest rate for a particular month is 6.6 percent. [3 marks]

Continued....

b) Alfred Industry produces electronic components that require four main raw materials. The table below shows the price and quantity required for each materials for the year of 2014 and 2015.

Raw material	2014		2015	
	Price (RM)	Quantity	Price (RM)	Quantity
A	3	40	4	35
B	5	27	5	32
C	4	25	6	30
D	6	10	7	20

Compute the Laspeyres Price Index (LPI) and Paashe Price Index (PPI) for 2015 using 2014 as the base year. [10 marks]

End of Question

A. DESCRIPTIVE STATISTICS

Mean $(\bar{x}) = \dfrac{\sum_{i=1}^{n} X_i}{n}$

Standard Deviation $(s) = \sqrt{\dfrac{\sum_{i=1}^{n} X_i^2}{n-1} - \dfrac{(\sum_{i=1}^{n} X_i)^2}{n(n-1)}}$

Coefficient of Variation $(CV) = \dfrac{\sigma}{\bar{X}} \times 100$

Pearson's Coefficient of Skewness $(S_k) = \dfrac{3(\bar{X} - Median)}{s}$

B. PROBABILITY

P (A or B) = P (A) + P (B) – P (A and B)

P (A and B) = P (A)× P (B) if A and B are independent

P (A | B) = P (A and B) ÷ P (B)

Poisson Probability Distribution

If X follows a Poisson Distribution , $P(\lambda)$ where $P(X = x) = \dfrac{e^{-\lambda}\lambda^x}{x!}$

then the mean = $E(X) = \lambda$ and variance = $VAR(X) = \lambda$

Binomial Probability Distribution

If X follows a Binomial Distribution B(n, p) where $P(X = x) = {}^nC_x p^x q^{n-x}$

then the mean = $E(X) = np$ and variance = $VAR(X) = npq$ where $q = 1 - p$

Normal Distribution

If X follows a Normal distribution, $N(\mu, \sigma)$ where $E(X) = \mu$ and $VAR(X) = \sigma^2$

then $Z = \dfrac{X - \mu}{\sigma}$

C. EXPECTATION AND VARIANCE OPERATORS

$E(X) = \sum[X \bullet P(X)]$

$VAR(X) = E(X^2) - [E(X)]^2$ where $E(X^2) = \sum[X^2 \bullet P(X)]$

If $E(X) = \mu$ then $E(cX) = c\mu$, $E(X_1 + X_2) = E(X_1) + E(X_2)$

If $VAR(X) = \sigma^2$ then $VAR(cX) = c^2\sigma^2$, $VAR(X_1 + X_2) = VAR(X_1) + VAR(X_2) + 2\,COV(X_1, X_2)$

where $COV(X_1, X_2) = E(X_1X_2) - [E(X_1)\,E(X_2)]$

D. CONFIDENCE INTERVAL ESTIMATION AND SAMPLE SIZE DETERMINATION

(100 – α) % Confidence Interval for Population Mean (σ Known) = $\mu = \overline{X} \pm Z_{\alpha/2}\left(\sigma/\sqrt{n}\right)$

(100 – α)% Confidence Interval for Population Mean (σ Unknown) = $\mu = \overline{X} \pm t_{\alpha/2,n-1}\left(S/\sqrt{n}\right)$

(100 – α)% Confidence Interval for Population Proportion $= \hat{p} \pm Z_{\alpha/2}\sigma_{\hat{p}}$ Where $\sigma_{\hat{p}} = \sqrt{\dfrac{\hat{p}(1-\hat{p})}{n}}$

Sample Size Determination for Population Mean $= n \geq \left[\dfrac{(Z_{\alpha/2})\sigma}{E}\right]^2$

Sample Size Determination for Population Proportion $= n \geq \dfrac{(Z_{\alpha/2})^2\,\hat{p}(1-\hat{p})}{E^2}$

Where E = Limit of Error in Estimation

E. HYPOTHESIS TESTING

One Sample Mean Test	
Standard Deviation (σ) Known	**Standard Deviation (σ) Not Known**
$Z = \dfrac{x-u}{\sigma/\sqrt{n}}$	$t = \dfrac{x-u}{s/\sqrt{n}}$
One Sample Proportion Test	
$z = \dfrac{\hat{p}-p}{\sigma_p}$ where $\sigma_p = \sqrt{\dfrac{p(1-p)}{n}}$	
Two Sample Mean Test	
Standard Deviation (σ) Known $z = \dfrac{(\overline{x}_1 - \overline{x}_2) - (\mu_1 - \mu_2)}{\sqrt{\sigma_1^2/n_1 + \sigma_2^2/n_2}}$	**Standard Deviation (σ) Not Known** $t = \dfrac{(\overline{x}_1 - \overline{x}_2) - (\mu_1 - \mu_2)}{\sqrt{S_p^2\left(1/n_1 + 1/n_2\right)}}$ where $S_p^2 = \dfrac{(n_1-1)S_1^2 + (n_2-1)S_2^2}{(n_1+n_2-2)}$
Two Sample Proportion Test	
$Z = \dfrac{\hat{p}_1 - \hat{p}_2}{\sqrt{p(1-p)\left(\frac{1}{n_1}+\frac{1}{n_2}\right)}}$ where $p = \dfrac{X_1 + X_2}{n_1 + n_2}$ *where X_1 and X_2 are the number of successes from each population*	

F. REGRESSION ANALYSIS

Simple Linear Regression

Population Model: $Y = \beta_0 + \beta_1 X_1 + \varepsilon$

Sample Model: $\hat{y} = b_0 + b_1 x_1 + e$

Correlation Coefficient

$$r = \frac{\sum XY - \left[\frac{\sum X \sum Y}{n}\right]}{\sqrt{\left[\sum X^2 - \left(\frac{(\sum X)^2}{n}\right)\right]\left[\sum Y^2 - \left(\frac{(\sum Y)^2}{n}\right)\right]}} = \frac{COV(X,Y)}{\sigma_X \sigma_Y}$$

ANOVA Table for Regression

Source	Degrees of Freedom	Sum of Squares	Mean Squares
Regression	*1*	SSR	MSR = SSR/1
Error/Residual	$n-2$	SSE	MSE = SSE/$(n-2)$
Total	$n-1$	SST	

Test Statistic for Significance of the Predictor Variable

$t_i = \frac{b_i}{S_{b_i}}$ and the critical value $= \pm t_{\alpha/2,(n-p-1)}$

Where p= number of predictor

G. INDEX NUMBERS

Simple Price Index $P = \frac{p_t}{p_0} \times 100$	**Laspeyres Quantity Index** $P = \frac{\sum p_0 q_t}{\sum p_0 q_0} \times 100$
Aggregate Price Index $P = \frac{\sum p_t}{\sum p_0}(100)$	**Paasche Quantity Index** $P = \frac{\sum p_t q_t}{\sum p_t q_0} \times 100$
Laspeyres Price Index $P = \frac{\sum p_t q_0}{\sum p_0 q_0} \times 100$	**Fisher's Ideal Price Index** $\sqrt{(Laspeyres\ Price\ Index)(Paa\ sche\ Price\ Index)}$
Paasche Price Index $P = \frac{\sum p_t q_t}{\sum p_0 q_t} \times 100$	**Value Index** $V = \frac{\sum p_t q_t}{\sum p_0 q_0} \times 100$

TABLE 4. THE NORMAL DISTRIBUTION FUNCTION

The function tabulated is $\Phi(x) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{x} e^{-\frac{1}{2}t^2}\, dt$. $\Phi(x)$ is the probability that a random variable, normally distributed with zero mean and unit variance, will be less than or equal to x. When $x < 0$ use $\Phi(x) = 1 - \Phi(-x)$, as the normal distribution with zero mean and unit variance is symmetric about zero.



x	$\Phi(x)$	x	$\Phi(x)$	x	$\Phi(x)$	x	$\Phi(x)$	x	$\Phi(x)$	x	$\Phi(x)$
0·00	0·5000	0·40	0·6554	0·80	0·7881	1·20	0·8849	1·60	0·9452	2·00	0·97725
·01	·5040	·41	·6591	·81	·7910	·21	·8869	·61	·9463	·01	·97778
·02	·5080	·42	·6628	·82	·7939	·22	·8888	·62	·9474	·02	·97831
·03	·5120	·43	·6664	·83	·7967	·23	·8907	·63	·9484	·03	·97882
·04	·5160	·44	·6700	·84	·7995	·24	·8925	·64	·9495	·04	·97932
0·05	0·5199	0·45	0·6736	0·85	0·8023	1·25	0·8944	1·65	0·9505	2·05	0·97982
·06	·5239	·46	·6772	·86	·8051	·26	·8962	·66	·9515	·06	·98030
·07	·5279	·47	·6808	·87	·8078	·27	·8980	·67	·9525	·07	·98077
·08	·5319	·48	·6844	·88	·8106	·28	·8997	·68	·9535	·08	·98124
·09	·5359	·49	·6879	·89	·8133	·29	·9015	·69	·9545	·09	·98169
0·10	0·5398	0·50	0·6915	0·90	0·8159	1·30	0·9032	1·70	0·9554	2·10	0·98214
·11	·5438	·51	·6950	·91	·8186	·31	·9049	·71	·9564	·11	·98257
·12	·5478	·52	·6985	·92	·8212	·32	·9066	·72	·9573	·12	·98300
·13	·5517	·53	·7019	·93	·8238	·33	·9082	·73	·9582	·13	·98341
·14	·5557	·54	·7054	·94	·8264	·34	·9099	·74	·9591	·14	·98382
0·15	0·5596	0·55	0·7088	0·95	0·8289	1·35	0·9115	1·75	0·9599	2·15	0·98422
·16	·5636	·56	·7123	·96	·8315	·36	·9131	·76	·9608	·16	·98461
·17	·5675	·57	·7157	·97	·8340	·37	·9147	·77	·9616	·17	·98500
·18	·5714	·58	·7190	·98	·8365	·38	·9162	·78	·9625	·18	·98537
·19	·5753	·59	·7224	·99	·8389	·39	·9177	·79	·9633	·19	·98574
0·20	0·5793	0·60	0·7257	1·00	0·8413	1·40	0·9192	1·80	0·9641	2·20	0·98610
·21	·5832	·61	·7291	·01	·8438	·41	·9207	·81	·9649	·21	·98645
·22	·5871	·62	·7324	·02	·8461	·42	·9222	·82	·9656	·22	·98679
·23	·5910	·63	·7357	·03	·8485	·43	·9236	·83	·9664	·23	·98713
·24	·5948	·64	·7389	·04	·8508	·44	·9251	·84	·9671	·24	·98745
0·25	0·5987	0·65	0·7422	1·05	0·8531	1·45	0·9265	1·85	0·9678	2·25	0·98778
·26	·6026	·66	·7454	·06	·8554	·46	·9279	·86	·9686	·26	·98809
·27	·6064	·67	·7486	·07	·8577	·47	·9292	·87	·9693	·27	·98840
·28	·6103	·68	·7517	·08	·8599	·48	·9306	·88	·9699	·28	·98870
·29	·6141	·69	·7549	·09	·8621	·49	·9319	·89	·9706	·29	·98899
0·30	0·6179	0·70	0·7580	1·10	0·8643	1·50	0·9332	1·90	0·9713	2·30	0·98928
·31	·6217	·71	·7611	·11	·8665	·51	·9345	·91	·9719	·31	·98956
·32	·6255	·72	·7642	·12	·8686	·52	·9357	·92	·9726	·32	·98983
·33	·6293	·73	·7673	·13	·8708	·53	·9370	·93	·9732	·33	·99010
·34	·6331	·74	·7704	·14	·8729	·54	·9382	·94	·9738	·34	·99036
0·35	0·6368	0·75	0·7734	1·15	0·8749	1·55	0·9394	1·95	0·9744	2·35	0·99061
·36	·6406	·76	·7764	·16	·8770	·56	·9406	·96	·9750	·36	·99086
·37	·6443	·77	·7794	·17	·8790	·57	·9418	·97	·9756	·37	·99111
·38	·6480	·78	·7823	·18	·8810	·58	·9429	·98	·9761	·38	·99134
·39	·6517	·79	·7852	·19	·8830	·59	·9441	·99	·9767	·39	·99158
0·40	0·6554	0·80	0·7881	1·20	0·8849	1·60	0·9452	2·00	0·9772	2·40	0·99180

TABLE 4. THE NORMAL DISTRIBUTION FUNCTION

x	$\Phi(x)$	x	$\Phi(x)$	x	$\Phi(x)$	x	$\Phi(x)$	x	$\Phi(x)$	x	$\Phi(x)$
2·40	0·99180	2·55	0·99461	2·70	0·99653	2·85	0·99781	3·00	0·99865	3·15	0·99918
·41	·99202	·56	·99477	·71	·99664	·86	·99788	·01	·99869	·16	·99921
·42	·99224	·57	·99492	·72	·99674	·87	·99795	·02	·99874	·17	·99924
·43	·99245	·58	·99506	·73	·99683	·88	·99801	·03	·99878	·18	·99926
·44	·99266	·59	·99520	·74	·99693	·89	·99807	·04	·99882	·19	·99929
2·45	0·99286	2·60	0·99534	2·75	0·99702	2·90	0·99813	3·05	0·99886	3·20	0·99931
·46	·99305	·61	·99547	·76	·99711	·91	·99819	·06	·99889	·21	·99934
·47	·99324	·62	·99560	·77	·99720	·92	·99825	·07	·99893	·22	·99936
·48	·99343	·63	·99573	·78	·99728	·93	·99831	·08	·99896	·23	·99938
·49	·99361	·64	·99585	·79	·99736	·94	·99836	·09	·99900	·24	·99940
2·50	0·99379	2·65	0·99598	2·80	0·99744	2·95	0·99841	3·10	0·99903	3·25	0·99942
·51	·99396	·66	·99609	·81	·99752	·96	·99846	·11	·99906	·26	·99944
·52	·99413	·67	·99621	·82	·99760	·97	·99851	·12	·99910	·27	·99946
·53	·99430	·68	·99632	·83	·99767	·98	·99856	·13	·99913	·28	·99948
·54	·99446	·69	·99643	·84	·99774	·99	·99861	·14	·99916	·29	·99950
2·55	0·99461	2·70	0·99653	2·85	0·99781	3·00	0·99865	3·15	0·99918	3·30	0·99952

The critical table below gives on the left the range of values of x for which $\Phi(x)$ takes the value on the right, correct to the last figure given; in critical cases, take the upper of the two values of $\Phi(x)$ indicated.

3·075	0·9990	3·263	0·9994	3·731	0·99990	3·916	0·99995
3·105	0·9991	3·320	0·9995	3·759	0·99991	3·976	0·99996
3·138	0·9992	3·389	0·9996	3·791	0·99992	4·055	0·99997
3·174	0·9993	3·480	0·9997	3·826	0·99993	4·173	0·99998
3·215	0·9994	3·615	0·9998	3·867	0·99994	4·417	0·99999
			0·9999		0·99995		1·00000

When $x > 3{\cdot}3$ the formula $1-\Phi(x) \doteqdot \frac{e^{-\frac{1}{2}x^2}}{x\sqrt{2\pi}}\left[1-\frac{1}{x^2}+\frac{3}{x^4}-\frac{15}{x^6}+\frac{105}{x^8}\right]$ is very accurate, with relative error less than $945/x^{10}$.

TABLE 5. PERCENTAGE POINTS OF THE NORMAL DISTRIBUTION

This table gives percentage points $x(P)$ defined by the equation

$$\frac{P}{100} = \frac{1}{\sqrt{2\pi}} \int_{x(P)}^{\infty} e^{-\frac{1}{2}t^2}\, dt.$$

If X is a variable, normally distributed with zero mean and unit variance, $P/100$ is the probability that $X \geqslant x(P)$. The lower P per cent points are given by symmetry as $-x(P)$, and the probability that $|X| \geqslant x(P)$ is $2P/100$.



P	$x(P)$	P	$x(P)$	P	$x(P)$	P	$x(P)$	P	$x(P)$	P	$x(P)$
50	0·0000	5·0	1·6449	3·0	1·8808	2·0	2·0537	1·0	2·3263	0·10	3·0902
45	0·1257	4·8	1·6646	2·9	1·8957	1·9	2·0749	0·9	2·3656	0·09	3·1214
40	0·2533	4·6	1·6849	2·8	1·9110	1·8	2·0969	0·8	2·4089	0·08	3·1559
35	0·3853	4·4	1·7060	2·7	1·9268	1·7	2·1201	0·7	2·4573	0·07	3·1947
30	0·5244	4·2	1·7279	2·6	1·9431	1·6	2·1444	0·6	2·5121	0·06	3·2389
25	0·6745	4·0	1·7507	2·5	1·9600	1·5	2·1701	0·5	2·5758	0·05	3·2905
20	0·8416	3·8	1·7744	2·4	1·9774	1·4	2·1973	0·4	2·6521	0·01	3·7190
15	1·0364	3·6	1·7991	2·3	1·9954	1·3	2·2262	0·3	2·7478	0·005	3·8906
10	1·2816	3·4	1·8250	2·2	2·0141	1·2	2·2571	0·2	2·8782	0·001	4·2649
5	1·6449	3·2	1·8522	2·1	2·0335	1·1	2·2904	0·1	3·0902	0·0005	4·4172

TABLE 10. PERCENTAGE POINTS OF THE t-DISTRIBUTION

This table gives percentage points $t_\nu(P)$ defined by the equation

$$\frac{P}{100} = \frac{1}{\sqrt{\nu\pi}} \frac{\Gamma(\frac{1}{2}\nu+\frac{1}{2})}{\Gamma(\frac{1}{2}\nu)} \int_{t_\nu(P)}^{\infty} \frac{dt}{(1+t^2/\nu)^{\frac{1}{2}(\nu+1)}}.$$

Let X_1 and X_2 be independent random variables having a normal distribution with zero mean and unit variance and a χ^2-distribution with ν degrees of freedom respectively; then $t = X_1/\sqrt{X_2/\nu}$ has Student's t-distribution with ν degrees of freedom, and the probability that $t \geqslant t_\nu(P)$ is $P/100$. The lower percentage points are given by symmetry as $-t_\nu(P)$, and the probability that $|t| \geqslant t_\nu(P)$ is $2P/100$.



The limiting distribution of t as ν tends to infinity is the normal distribution with zero mean and unit variance. When ν is large interpolation in ν should be harmonic.

P	40	30	25	20	15	10	5	2·5	1	0·5	0·1	0·05
$\nu = 1$	0·3249	0·7265	1·0000	1·3764	1·963	3·078	6·314	12·71	31·82	63·66	318·3	636·6
2	0·2887	0·6172	0·8165	1·0607	1·386	1·886	2·920	4·303	6·965	9·925	22·33	31·60
3	0·2767	0·5844	0·7649	0·9785	1·250	1·638	2·353	3·182	4·541	5·841	10·21	12·92
4	0·2707	0·5686	0·7407	0·9410	1·190	1·533	2·132	2·776	3·747	4·604	7·173	8·610
5	0·2672	0·5594	0·7267	0·9195	1·156	1·476	2·015	2·571	3·365	4·032	5·893	6·869
6	0·2648	0·5534	0·7176	0·9057	1·134	1·440	1·943	2·447	3·143	3·707	5·208	5·959
7	0·2632	0·5491	0·7111	0·8960	1·119	1·415	1·895	2·365	2·998	3·499	4·785	5·408
8	0·2619	0·5459	0·7064	0·8889	1·108	1·397	1·860	2·306	2·896	3·355	4·501	5·041
9	0·2610	0·5435	0·7027	0·8834	1·100	1·383	1·833	2·262	2·821	3·250	4·297	4·781
10	0·2602	0·5415	0·6998	0·8791	1·093	1·372	1·812	2·228	2·764	3·169	4·144	4·587
11	0·2596	0·5399	0·6974	0·8755	1·088	1·363	1·796	2·201	2·718	3·106	4·025	4·437
12	0·2590	0·5386	0·6955	0·8726	1·083	1·356	1·782	2·179	2·681	3·055	3·930	4·318
13	0·2586	0·5375	0·6938	0·8702	1·079	1·350	1·771	2·160	2·650	3·012	3·852	4·221
14	0·2582	0·5366	0·6924	0·8681	1·076	1·345	1·761	2·145	2·624	2·977	3·787	4·140
15	0·2579	0·5357	0·6912	0·8662	1·074	1·341	1·753	2·131	2·602	2·947	3·733	4·073
16	0·2576	0·5350	0·6901	0·8647	1·071	1·337	1·746	2·120	2·583	2·921	3·686	4·015
17	0·2573	0·5344	0·6892	0·8633	1·069	1·333	1·740	2·110	2·567	2·898	3·646	3·965
18	0·2571	0·5338	0·6884	0·8620	1·067	1·330	1·734	2·101	2·552	2·878	3·610	3·922
19	0·2569	0·5333	0·6876	0·8610	1·066	1·328	1·729	2·093	2·539	2·861	3·579	3·883
20	0·2567	0·5329	0·6870	0·8600	1·064	1·325	1·725	2·086	2·528	2·845	3·552	3·850
21	0·2566	0·5325	0·6864	0·8591	1·063	1·323	1·721	2·080	2·518	2·831	3·527	3·819
22	0·2564	0·5321	0·6858	0·8583	1·061	1·321	1·717	2·074	2·508	2·819	3·505	3·792
23	0·2563	0·5317	0·6853	0·8575	1·060	1·319	1·714	2·069	2·500	2·807	3·485	3·768
24	0·2562	0·5314	0·6848	0·8569	1·059	1·318	1·711	2·064	2·492	2·797	3·467	3·745
25	0·2561	0·5312	0·6844	0·8562	1·058	1·316	1·708	2·060	2·485	2·787	3·450	3·725
26	0·2560	0·5309	0·6840	0·8557	1·058	1·315	1·706	2·056	2·479	2·779	3·435	3·707
27	0·2559	0·5306	0·6837	0·8551	1·057	1·314	1·703	2·052	2·473	2·771	3·421	3·690
28	0·2558	0·5304	0·6834	0·8546	1·056	1·313	1·701	2·048	2·467	2·763	3·408	3·674
29	0·2557	0·5302	0·6830	0·8542	1·055	1·311	1·699	2·045	2·462	2·756	3·396	3·659
30	0·2556	0·5300	0·6828	0·8538	1·055	1·310	1·697	2·042	2·457	2·750	3·385	3·646
32	0·2555	0·5297	0·6822	0·8530	1·054	1·309	1·694	2·037	2·449	2·738	3·365	3·622
34	0·2553	0·5294	0·6818	0·8523	1·052	1·307	1·691	2·032	2·441	2·728	3·348	3·601
36	0·2552	0·5291	0·6814	0·8517	1·052	1·306	1·688	2·028	2·434	2·719	3·333	3·582
38	0·2551	0·5288	0·6810	0·8512	1·051	1·304	1·686	2·024	2·429	2·712	3·319	3·566
40	0·2550	0·5286	0·6807	0·8507	1·050	1·303	1·684	2·021	2·423	2·704	3·307	3·551
50	0·2547	0·5278	0·6794	0·8489	1·047	1·299	1·676	2·009	2·403	2·678	3·261	3·496
60	0·2545	0·5272	0·6786	0·8477	1·045	1·296	1·671	2·000	2·390	2·660	3·232	3·460
120	0·2539	0·5258	0·6765	0·8446	1·041	1·289	1·658	1·980	2·358	2·617	3·160	3·373
∞	0·2533	0·5244	0·6745	0·8416	1·036	1·282	1·645	1·960	2·326	2·576	3·090	3·291